

X-D. Experimental Atherosclerosis

In the course of investigating the toxicity of nicotine, Adler and Hensel (1906) discovered lesions in the aorta of rabbits. The lesions were degenerative and aneurysmal; they were similar to those induced by epinephrine and unlike the human form of atherosclerosis. Degenerative lesions were later seen in rabbits by Grosogeat and Roubelakis (1965) and Lellouch et al. (1968) and in dogs by Hueper (1943) but were not seen in rats by Thienes (1960). The chronic administration of nicotine alone caused a transient increase in blood lipids but the level approximated the control level towards the end of the experiment (Schivelbein et al., 1970). Although the calcium controls of the aorta of nicotine-treated rabbits was increased, there were no differences in histological changes in the blood vessels between nicotine-treated and control rabbits.

The rabbit has been the favorite animal for experimental induction of atherosclerosis. The initial attempt of Wenzel et al. (1959) failed to establish an increase in severity of atherosclerosis in rabbits on a high cholesterol diet. Two other groups reported an increase in extent of dietary-induced atherosclerosis in nicotine-treated rabbits—one group from Chicago (Haas et al., 1966a, 1966b, 1968; Landerholm et al., 1967) and another group from Boston (Stefanovich et al., 1969). The oxygen uptake of tissues from hypercholesterolemic rabbits has been reported to be more sensitive to the depressant action of nicotine than that of tissues from

1005050490

normocholesterolemic rabbits (Adachi et al., 1965). Nicotine was shown to change the lipid patterns of the perfused dog aorta and coronary artery (Kupke, 1972). The results were regarded by the author as supporting the hypothesis that nicotine may impair oxidative enzymes by damaging the mitochondrial structures, thereby leading to lipid accumulation. Experiments on the atherosclerotic dog aorta and the human aorta are needed.

1005050491

BIBLIOGRAPHY

CIRCULATORY EFFECTS OF NICOTINE

D. Experimental Atherosclerosis

	Reprint
ADACHI T, ADACHI M and POLLAK O J: Nicotine effect on oxygen consumption of tissues from normal and hypercholesteremic rabbits. <u>Fed Proc</u> 24: 721, 1965.	945
ADLER I and HENSEL O: Intravenous injections of nicotine and their effects upon the aorta of rabbits. <u>J Med Res</u> 10: 229-239, 1906.	946
GROSGOGÉAT Y and ROUBELAKIS G: Action expérimentale de la nicotine sur la paroi aortique du lapin. Étude en microscopie optique et électronique. <u>Pathol Biol</u> 13: 1140-1155, 1965.	947
HASS G M, LANDERHOLM W and HEMMENS A: Production of calcific athero-arteriosclerosis and thromboarteritis with nicotine, Vitamin D and dietary cholesterol. <u>Am J Pathol</u> 49: 739-741, 1966a.	948
HASS G M, LANDERHOLM W and HEMMENS A: Nicotine induction of arteriosclerosis and thromboarteritis in rabbits. <u>Circulation</u> 34 (Suppl 3): 14, 1966b.	949
HASS G M, HENSON D, LANDERHOLM W and HEMMENS A: Prevention of nicotine induction of atherocalcific thromboarteritis in rabbits. <u>Circulation</u> 38 (Suppl 6): 8, 1968.	950
HUEPER W C: Experimental studies in cardiovascular pathology. VII. Chronic nicotine poisoning in rats and in dogs. <u>Arch Pathol</u> 35: 846-856, 1943.	951
KUPKE I R: Biosynthesis of lipids in perfused dog aorta and coronary artery. <u>J Mol Cell Cardiol</u> 4: 27-38, 1972.	952
LANDERHOLM W, HEMMENS A and HASS G M: An experimental inquiry into nicotine-induced thromboarteritis. <u>Fed Proc</u> 26: 359, 1967.	953
LELLOUCH J, JACOTOT B, ANGUERA G, GROSGOGÉAT J and BEAUMONT J-L: Action chronique de la nicotine sur l'intima aortique du lapin. Influence d'un inhibiteur de la monoamine oxydase. <u>J Atherosclerosis Res</u> 8: 137-142, 1968.	954
SCHIEVELBEIN H, LONDONG V, LONDONG W, GRUMBACH H, REMPLIK V, SCHAUER A and IMMICH H: Nicotine and arteriosclerosis. <u>Z Clin Chem Klin Biochem</u> 8: 190-196, 1970.	955
STEFANOVICH V, GORE I, KAJIYAMA G and IWANAGA Y: The effect of nicotine on dietary atherogenesis in rabbits. <u>Exp Mol Pathol</u> 11: 71-81, 1969.	956
THIENES C H: Chronic nicotine poisoning. <u>Ann N Y Acad Sci</u> 90: 239-248, 1960.	957
WENZEL D G, TURNER J A and KISSIL D: Effect of nicotine on cholesterol-induced atherosclerosis in the rabbit. <u>Circ Res</u> 7: 256-261, 1959.	958

1005050492

X-E. Experimental Hyperlipidemias

Epinephrine is known to mobilize free fatty acids from fatty tissue and this was followed by elevation of cholesterol and triglycerides in the blood (Kaplan et al., 1957; Sussman et al., 1958). A similar response was noted to nicotine administration in dogs (White et al., 1964; Kershbaum et al., 1965, 1967a, 1967b). There is also an accompanying elevation of blood glucose related to the release of epinephrine which has a glycogenolytic action (Tsujimoto et al., 1965; Milton, 1966). Hyperlipidemia induced in dogs by nicotine can be prevented by prior injection of adrenergic blocking drugs (Kershbaum et al., 1966). Animals with hypercholesterolemia show an interaction with nicotine. The administration of nicotine and ergonovine produces necrosis in the heart (Wenzel et al., 1961). Grosogeat did not find lesions in the rabbit aortic wall with nicotine treatment alone (Grosogeat et al., 1965). There is a diminution in the rate of synthesis of cholesterol in dogs that have been subjected to chronic administration of nicotine (Gudbjarnason, 1968). The relationship of cholesterol to atherosclerotic lesions is discussed in the preceding section.

1005050493

BIBLIOGRAPHY

X. CIRCULATORY EFFECTS OF NICOTINE

E. Experimental Hyperlipidemias

Reprint

- GROSGOGGAT Y, ANGUERA G, LELLOUCH J, JACOTOT B and BEAUMONT J L: Intoxication chronique par la nicotine chez le lapin nourri au cholestérol. J Atherosclerosis Res 5: 291-301, 1965. 959
- GUDBJARNASON S: Effect of chronic nicotine administration on cholesterol metabolism of liver, serum, heart and brain. J Pharmacol Exp Ther 161: 47-54, 1968. 960
- KAPLAN A, JACQUES S and GANT M: Effect of long-lasting epinephrine on serum lipid levels. Am J Physiol 191: 8-12, 1957. 961
- KERSHBAUM A, BELLET S and KHORSANDIAN R: Elevation of serum cholesterol after administration of nicotine. Am Heart J 69: 206-210, 1965. 962
- KERSHBAUM A, BELLET S, HIRABAYASHI M and FEINBERG L J: Regular, filter-tip, and modified cigarettes. JAMA 201: 545-546, 1967a. 963
- KERSHBAUM A, OSADA H, SCIABINE A, BELLET S and PAPPAJOHN J: Influence of nicotine on the mobilization of free fatty acids from rat adipose tissue in vitro and in the isolated perfused dog limb. Circulation 36 (Suppl 2): 20, 1967b. 964
- KERSHBAUM A, JIMINEZ J, BELLET S and ZANUTTINI D: Modification of nicotine-induced hyperlipidemia by antiadrenergic agents. J Atherosclerosis Res 6: 524-530, 1966. 965
- MILTON A S: The effect of nicotine on blood glucose levels and plasma non-esterified fatty acid levels in the intact and adrenalectomized cat. Br J Pharmacol 26: 256-263, 1966. 966
- SUSSMAN K E, SHAFRIR E and STEINBERG D: Mobilization of lipids by epinephrine. Circulation 18: 486, 1958. 967
- TSUJIMOTO A, TANINO S and KUROGOCHI Y: Effect of nicotine on serum potassium and blood glucose. Jap J Pharmacol 15: 415-422, 1965. 968
- WENZEL D G, TURNER J A, JORDAN S W and SINGH J: Cardiovascular interaction of nicotine, ergonovine, and hypercholesterolemia in the rabbit. Circ Res 9: 694-699, 1961. 969
- WHITE H J, GORE I and LARKEY B J: The antagonism between nicotine and mucopolysaccharide activity. Biochim Biol Experimentale 3: 107-116, 1964. 970

1005050494

X-F. Experimental Thrombosis

The release of epinephrine from the adrenal medulla has been known to increase the coagulability of the blood in vitro (Cannon and Gray, 1914; Cannon and Mendenhall, 1914a, 1914b). More recently, this phenomenon regarding epinephrine has been demonstrated by techniques involving experimental thrombosis (Rosewell et al., 1966) and platelet aggregation (Shimamoto and Ishioka, 1963; Ardlie et al., 1966; Hampton and Mitchell, 1966; Besterman et al., 1967).

That nicotine releases epinephrine does not necessarily mean that all its reported effects on blood coagulation are mediated through this mechanism. Platelet aggregation results from the in vitro addition of nicotine, which must a significant amount of act directly on the cells not containing epinephrine (Werle and Schievelbein, 1965). In the rat, the thrombosis induced by a single subcutaneous injection of carragenine and nicotine can be simulated by substituting epinephrine for the nicotine (Jan et al., 1969). Chronic injection of nicotine in the same animal species caused decreased thrombus formation, indicating that there is no experimental support for the statement that habitual smokers are susceptible to thrombus formation (Wenzel and Richards, 1970). In rats maintained on a hypercholesterolemic diet, the administration of nicotine shortens coagulation time (Singh, 1965).

In other species, nicotine has varied effects, such as: reduction of toxicity of nicotine injected intravenously by increasing the number of platelets by transfusion in rabbit (Schievelbein and Schirren, 1964), decrease of

1005050495

fibrinolysin in the guinea pig by chronic administration of nicotine (Belli et al., 1965), and nicotine-heparin antagonism when added to human blood in vitro (Singh and Oester, 1964a, 1964b). These observations suggest that nicotine influences coagulability by several mechanisms other than the release of epinephrine.

1005050496

BIBLIOGRAPHY

X. CIRCULATORY EFFECTS OF NICOTINE

F. Experimental Thrombosis

Reprint

- ARDLIE N G, GLEW G and SCHWARTZ C J: Influence of catecholamines on nucleotide-induced platelet aggregation. Nature 212: 415-417, 1966. 971
- BELLI G, ALBERTINI E, MARTINESI L and PELLEGRINI G F: Studio sperimentale di momenti dell'emocoagulazione e della fibrinolisi in corso d'intossicazione nicotinic cronica. Arch Sci Med 120: 460-478, 1965. 972
- BESTERMAN E, MYAT G and TRAVADI V: Diurnal variations of platelet stickiness compared with effects produced by adrenaline. Br Med J 1: 597-600, 1967. 973
- CANNON W B and GRAY H: Factors affecting the coagulation time of blood. II. The hastening or retarding of coagulation by adrenalin injections. Am J Physiol 34: 232-242, 1914. 974
- CANNON W B and MENDENHALL W L: Factors affecting the coagulation time of blood. III. The hastening of coagulation by stimulating the splanchnic nerves. Am J Physiol 34: 243-250, 1914a. 975
- CANNON W B and MENDENHALL W L: Factors affecting the coagulation time of blood. IV. The hastening of coagulation in pain and emotional excitement. Am J Physiol 34: 251-261, 1914b. 976
- JAN F, LACOTOT B and BEAUMONT J-L: Phénomène thrombo-hémorragique déclenché par la nicotine. Fathol Biol 17: 501-504, 1969. 977
- HAMPTON J R and MITCHELL J R A: Effect of aggregating agents on the electrophoretic mobility of human platelets. Br Med J 1: 1074-1077, 1966. 978
- ROWSSELL H C, HEGARDT B, DOWNIE H G, MUSTARD J F and MURPHY E A: Adrenaline and experimental thrombosis. Br J Haematol 12: 66-73, 1966. 979
- SCHIEVELBEIN H and SCHIRREN V: Abschwächung der Toxizität von Nicotin durch Erhöhung der Thrombocytenzahl. Experienta 20: 432-433, 1964. 980
- SHIMAMOTO T and ISHIOKA T: Release of a thromboplastic substance from arterial walls by epinephrine. Circ Res 12: 138-144, 1963. 981
- SINGH J and OESTER Y T: Nicotine antagonism with heparin possible mode of action on human blood coagulation time in vitro. Arch Int Pharmacodyn 149: 354-361, 1964a. 982
- SINGH J and OESTER Y T: Effect of nicotine on prothrombin time and its possible mode of action. Arch Int Pharmacodyn 150: 435-441, 1964b. 983

1005050497

X. Acute and Chronic Effects
of Nicotine - 32


Page 285

Reprint

- SINGH J: Effect of nicotine on blood clotting time of albino rats fed atherogenic diet. Arch Int Pharmacodyn 154: 221-227, 1965. 984
- WENZEL D G and RICHARDS M H: Decreased thrombus formation in rats after chronic nicotine administration. Eur J Pharmacol 10: 143-144, 1970. 985
- WERLE E and SCHIEVELBEIN H: Activity of nicotine and inactivity of kaffikrein and kallidin in aggregation of blood platelets. Nature 207: 871-872, 1965. 986

1005050498

XI. CIRCULATORY EFFECTS OF CARBON MONOXIDE



X CARBON MONOXIDE

A. Carboxyhemoglobin Blood Levels-----	287
B. Influence of Smoking on Alveolar Air and Blood Levels of Carboxyhemoglobin-----	295
C. Coronary Circulation and Myocardium-----	300
D. Experimental Atherosclerosis--	308
E. Blood Cells and Plasma-----	315

The most important single measurement for assessing the extent of exposure to carbon monoxide contained in cigarette smoke is that of the blood level of carboxyhemoglobin. This form of hemoglobin reduces the availability of oxygen, so that there is a danger of producing generalized hypoxia. It has been suggested that the hypoxia in the myocardium would lead to coronary heart disease and vascular hypoxia/^{to} atherosclerosis. The experiments which form the basis of this statement are analyzed in the present section.

The author has completed a separate review entitled "Cigarette Smoking and Carbon Monoxide" and submitted it to the Council for Tobacco Research on January 3, 1973. It contains a discussion of the effects of carbon monoxide as it relates to cigarette smoking. The circulatory effects are recounted in this section and the literature has been updated to July 1, 1973.

1005050499

XI-A. Carboxyhemoglobin Blood Levels

The carboxyhemoglobin present in the blood is not able to dissociate the readily to allow/hemoglobin portion to combine with oxygen. The blood levels are reported for the habitual smoker compared with / the nonsmoker, and the acute elevation during cigarette smoking.

1. Habitual smokers compared / with nonsmokers. The first identification of carboxyhemoglobin in blood of smokers was accomplished by carboxyhemoglobin Hartridge (1919-1920). He estimated the/level in one smoker to be 6 % and in another to be absent. A more extensive investigation was completed by Gettler and Mattice (1933), who compared four groups of habitual cigarette smokers. The group of 12 rural dwellers had a mean value of 1.2 % carboxyhemoglobin in the blood, while 18 New York City residents had a mean of 1.4 %, 12 New York City cleaners a mean of 3.5 %, and two New York City taxi drivers a mean of 13.5 %. These results proved for the first time that the carboxyhemoglobin contained in blood of habitual smokers was not only the result of cigarette smoking but was also the outcome of inhaling an atmosphere containing carbon monoxide released from automobile exhaust and other sources.

There have been 28 additional reports of carboxyhemoglobin in blood levels among cigarette smokers and the results are summarized in Table XI-A. The mean values do not represent the effect of cigarette smoking because the contribution of carbon monoxide in the atmosphere has to be subtracted. The last column in Table XI-A is the net level of carboxyhemoglobin which can be attributed to cigarette smoking and was obtained by subtracting the mean

1005050500

blood levels for controls who were nonsmokers. The following generalizations can be made from the published results.

a. The 30 investigations summarized in Table XI-A were performed in various cities in the United States and Europe. The highest mean level is 16.2% for a group of 6 U. S. Army enlisted men (Meigs, 1948). This represents an error in the analysis, since the blood levels for nonsmokers were also high. All the other mean levels were below 10 % carboxyhemoglobin.

b. The overall mean level for 2,054 subjects reported in the 30 investigations is 3.76 %. This represents the average blood level for smokers in the morning, 4 to 12 hours after they smoked a cigarette.

c. The contribution of atmospheric pollution to the increased blood levels of carboxyhemoglobin can be derived by subtracting the blood levels for controls who were nonsmokers. Twenty-one of the investigations included nonsmokers, so that it was possible to subtract their mean levels from those of habitual smokers. The net difference between 2 groups represents the contribution from smoking alone, which amounted to a mean of +2.19% of carboxyhemoglobin for 2,781 subjects. This mean value was calculated regardless of the number of cigarettes consumed.

d. Ten investigators related the amount of cigarettes consumed daily to blood levels of carboxyhemoglobin. The consumption of 20 or less cigarettes per day showed the following net change in blood carboxyhemoglobin levels in each case: +1.6 % (Schmidt, 1939); +2.4% (Schrenk, 1942); +2.1% (Parmeggiani and Gilardi, 1952); +1.0% (Goldsmith et al., 1963); +1.9 %

1005050501

(Curphey et al., 1965) +0.3% (Balbo et al., 1966); and +0.9% (Rouch et al., 1971). The net changes in blood carboxyhemoglobin levels for subjects consuming one or more cigarettes were respectively: +2.9% (Schmidt, 1939); +3.7% (Schrenk, 1942); +3.5% (Parmeggiani and Gilardi, 1952); +1.1% (Goldsmith et al., 1963); +3.6% (Curphey et al., 1965); +2.0% (Balbo et al., 1966); and +8.0 (Yacoub et al., 1970). The last-mentioned value represents the highest net level of carboxyhemoglobin, next to the +11.0% referred to above as reported for the U. S. Army enlisted men.

(Table XI-A appears on the next page.)

1005050502

Table XI-A. Carboxyhemoglobin levels in the blood of habitual smokers.¹

Reference (year)	Nature of habitual smokers (Cigarette consumption)	No. of Subjects	Carboxyhemoglobin blood level %		
			Smokers Mean \pm SD (Range)	Non-smokers Mean	Net
Hartridge (1919-1920)	London volunteers	2	3.0 (0-6)		
Gottler and Mattice (1933)	New York residents	18	1.4 (1.0-4.1)		
	New York street cleaners	12	3.5 (1.2-0.9)		
	New York taxi drivers	2	13.5 (8.0-19)		
	Rural dwellers	12	1.2 (0.5-3.6)		
Ruhl and Lin (1936)	Berlin volunteers				
	(non-inhalers in morning)	21	0.6		
	(inhalers in morning)	25	0.5		
	(heavy inhalers in morning)	13	1.3		
Schmidt (1939; 1940)	Bonn volunteers				
	(20-30/day)	3	3.5	0.6	+2.9
	(>30/day)		7.2		+6.6
	(10-20/day)		2.2		+1.6
Schrenk (1942); Sievers, Edwards Murray and Schrenk (1942)	Holland tunnel workers				
	(<20/day)	39	4.1 \pm 1.9	1.7	+2.4
	(>20/day)	21	5.4 \pm 1.8		+3.7
	(pipe)	5	2.5		
Wennesland 1945	Stockholm volunteers				
	(<15/day)	35	(0.5-10.5)		
Meigs 1948	US army enlisted men				
	(6-20/day)	6	16.2 (1.9-45)	5.2	+11.0
Parmeggiani and Gibaldi (1952)	Italian volunteers				
	(10-12/day)	14	4.9 (1-9)	2.8	+2.1
	(15-25/day)	6	6.3 (2-9.5)		+3.5
Barthe, Paris, Duchemin and Thomas (1953)	Paris workers				
	(30-40/day)	3	9.3 (6-14)		+6.5
		100	(0.8-2.0)		
Ruel and Barthe (1954)	Paris workers				
	(<10/day)		1.0		
	(10-15/day)		1.7		
Valic and Duric (1954)	Yugoslavian workers				
	(>20/day)		2.7		
		75	3.8 \pm 1.9	0.9	+2.9
Cler, Cadigan, Wilcott, Jones and Mark (1957)	Boston workers				
		9	4.7 (3.1-7.9)	0.9	+3.8
Dahlström, Nordström, Örberg and Rothschild (1958)	Stockholm volunteers				
		6	2.1 (1.2-3.0)	0.7	+1.4

1005050503

Reference (Year)	Nature of habitual smokers (Cigarette consumption)	No. of Subjects	Carboxyhemoglobin blood level %			
			Smokers Mean \pm SD (Range)	Non-smokers ² Mean	No.	
Sunder and Harper (1962)	Sunderland volunteers	3	3.8 (3.2-4.8)	0.9	+2.9	
Hofreuter, Catcott and Xintaras (1962)	Cincinnati volunteers	19	2.9	1.9	+1.0	
Goldsmith, Schuette and Novick (1963)	San Francisco longshoremen (<10 /day)	429	2.3	1.3	+1.0	
	(10-40/day)	1035	3.4		+1.1	
	(>40 /day)	233	5.5		+4.2	
Curphrey, Hood and Perkins (1965)	Los Angeles longshoremen (light)	55	2.3	0.4	+1.9	
	(medium)	153	3.0		+2.0	
	(heavy)	29	4.0		+3.6	
Ayres, Gianelli and Armstrong (1965)	New York volunteers	25	4.2	0.9	+3.3	
Balbo, Marucci and Ronchi (1966)	Paris workers (5/day)	34	2.1 (0-9.8)	1.9	+0.2	
	(10/day)	32	2.2 (0-9.5)		+0.3	
	(15/day)	16	2.2 (0-5.9)		+0.3	
	(20/day)	20	2.2 (0-6.9)		+0.3	
	(30/day)	7	2.8 (0-5.2)		+2.0	
Quivaine, Nelson and Bartlett (1969)	Durham workers	5	3.8	1.6	+2.2	
Bhown, Maitrya and Haq (1969)	Indian beedi smokers (<10 /day)	7	4.8 (3.4-5.6)			
	(10-19/day)	8	5.9 (5.6-6.1)			
	(20-29/day)	8	6.9 (6.6-7.3)			
	(>29 /day)	7	9.4 (8.5-10.5)			
Yacoub, Faure, Mallion and Cau (1970)	Paris workers (20/day inhaled)	90	9.5	1.5	+8.0	
	(20/day non-inhaled)	97	6.0		+4.5	
Rouch, Rioufol and Bourbon (1971)	Toulouse volunteers (<10 /day)	5	2.5 (1.0-5.5)	1.6	+0.9	
	(>10 /day)	15	4.25 (1.0-11.0)		+2.6	
Brewer, Eaton, Weil and Grover (1970); Brewer, Eaton, Grover and Weil (1971)	Leadville volunteers	20	6.6 \pm 2.7			
Weiss, Slawsky and Desforges (1971)	Boston patients with fibrosis	8	4.2 \pm 2.7	1.5	+4.2	
Seely (1971)	Los Angeles residents	81	5.7 (3.2-14.2)	1.5	+4.2	
Hansen, Wilke, Malbrny and Gøthert (1972)	Hamburg workers	40	4.9 \pm 1.0	0.75	+4.2	

1005050504

Reference (Year)	Nature of habitual smokers (Cigarette consumption)	No. of Subjects	Carboxyhemoglobin blood levels%		
			Smokers Mean \pm SD (Range)	Non-smokers Mean	Net
Person, Garby, bert and Zaar (1972)	Uppsala volunteers	6	2.7 (1.5-4.4)	1.27	+1.4
Mean (overall for number of subjects)			3.76 (2954 subjects)		+2.19 (2781 subjects)

¹ Some of the values were reported in volumes % and are expressed in this table as saturation % assuming normal hemoglobin values.

² See Table 4 for details.

1005050505

BIBLIOGRAPHY

XI. CIRCULATORY EFFECTS OF CARBON MONOXIDE

B. Influence of Smoking on Alveolar Air and Blood Levels of Carboxyhemoglobin

- BARACH A L, ECKMAN M and MOLOMUT N: Modification of resistance to anoxia with especial reference to high altitude flying. Am J Med Sci 202: 336-41, 1941. 1018
- BELLI R and GUILLANO V: Ricerche sull'ossicarbonismo. Ossicarbonemia da fumo di tabacco e diagnosi di ossicarbonismo cronico. Foll Med (Napoli) 38: 351-8, 1955. 1019
- BOWDEN C H and WOODHALL W R: The determination and significance of low blood carboxyhaemoglobin levels. Med Sci Law 4: 98-107, 1964. 1020
- CASTELLINO N: Modifiche della carbossiemia in seguito alla inalazione del fumo di tabacco. Foll Med (Napoli) 38: 1014-24, 1955. 1021
- COHEN S I, PERKINS N M, URY H K and GOLDSMITH J R: Carbon monoxide uptake in cigarette smoking. Arch Environ Health 22: 55-60, 1971. 1022
- CROSETTI L, RUBINO G F and PETTINATI L: Osservazioni in tema di detossicazione dell'emoglobina nell'uomo esposto a rischio prolungato da ossido di carbonio. Minerva Med 57: 268-9, 1966. 1023
- DONTENWILL W, RECKZEH G and STADLER L: Inhalationsexperimente mit Zigarettenrauch. Beitr Tabakforsch 3: 438-48, 1966. 1024
- DONTENWILL W, RECKZEH G and STADLER L: Berauchungsapparatur für Laboratoriumstiere. (Smoking apparatus for laboratory animals). Beitr Tabakforsch 4: 45-9, 1967. 1025
- DRISCOLL P, DEUBER A, BAETTIG K and GRANDJEAN E: Effect of filtered cigarette smoke on rats. Nature 237: 37-8, 1972. 1026
- FABRE R, TRUHAUT R and BERROD F: Une nouvelle méthode de dosage de l'oxyde de carbone dans le sang. Applications à l'étude de l'oxycarbonémie normale et pathologique. Ann Pharm (France) 9: 625-38, 1951. 1027
- and HACKNEY J D
- GOLDSMITH J R, TERZAGHI J: Evaluation of fluctuating carbon monoxide exposures. Arch Environ Health 7: 647-63, 1963. 1028
- HAMILL W and O'NEILL R P: Carbon monoxide intoxication in cigar smokers. Irish J Med Sci 8: 273-7, 1969. 1029
- HANSON H B and HASTINGS A B: The effect of smoking on the carbon monoxide content of blood. JAMA 100: 1481, 1933. 1030
- HSI-PU S and LI-MING: Carbon monoxide content of the blood after smoking. J Oriental Med 32: 1135-9, 1940. 1031
- JONGBLOED J: Über den Kohlenoxydgehalt der Alveolarluft beim Tabakrauchen. Arch Int Pharmacodyn 63: 346-58, 1939. 1032
- LEUCHTENBERGER C, LEUCHTENBERGER R and WEISS S: Increase of carbon monoxide (CO) in blood of mice exposed to inhalation of cigarette smoke. Proc Am Ass Canc Res 6: 39, 1965. 1033
- MacFARLAND R A, ROUGHTON F J W, HALPERIN M H and NIVEN J L: The effects of carbon monoxide and altitude on visual thresholds. J Aviation Med 15: 381- , 1944. 1034
- PARMEGGIANI L and GILARDI F: Rilievi sulla ossicarbonemia fisiologica. Med Lav 43: 179-83, 1952. 1035
- RAMSEY A: Fall in carbon monoxide blood levels after stopping smoking. Lancet I: 755, 1977. 1036

1005050506

- RINGOLD A, GOLDSMITH J R, HELWIG H I, FINN R and SCHUETTE F: Estimating recent carbon monoxide exposures. A rapid method. Arch Environ Health 5: 308-18, 1962. 1037
- ROSENBERG A: Kontrolleret tobakafvaenning. (Controlled tobacco detoxication). Ugeskrift for Laeger 130: 2014, 1968. 1038
- ROSENBERG A: Synpunkter pa rokavvanjning - erfarenheter fran Kopenhamn. (Viewpoints on smoking withdrawal - Copenhagen experiences. Soc-Med Tidskrifts 2: 103-10, 1971). 1039
- ROSENBERG A: Fall in carbon monoxide blood levels after stopping smoking. Lancet 1: 593, 1972. 1040
- RUHL A and LIN P: Zur frage der Kohlenoxydintoxikation bei starken Rauchern. Deutsch Med Wschr 62: 493-7, 1936. 1041
- SCHMIDT O: Der gasanalytische nachweis von Kohlenoxyd im Blut, insbesondere beim Rauchern. Klin Wschr 18: 938, 1939. 1042
- SCHMIDT O: Der Kohlenoxydgehalt des Blutes bei Rauchern. (The CO content of the blood of smokers). Reichs-GesundHbl 15: 53-8, 1940. 1043
- SHIELDS C E: Elevated carbon monoxide levels from smoking in blood donors. Transfusion 11: 89-93, 1971. 1044
- WHITEHEAD T P and WORTHINGTON S: The determination of carboxyhaemoglobin. Clin Chim Acta 6: 356-9, 1961. 1045

1005050507